

## Comparison between Sesame Oil and Glycyrrhizaglabra Effect as Phytoestrogen on Male Albino Rats

Eman G.E. Helal<sup>1</sup>, Mohamed A. El-Aleem<sup>2</sup>, Shaimaa S. Ahmed<sup>1</sup>

Zoology Department, Faculty of Science, Al-Azhar University (Girls), Cairo, Egypt.<sup>1</sup>

Physiology Department, Faculty of Medicine, Al-Azhar University, Cairo, Egypt<sup>2</sup>

\*Corresponding Author: Eman G.E. Helal, E-mail: emanhelal@hotmail.com, Mobile: 00201001025364,  
Orchid.org/0000-0003-0527-7028

### ABSTRACT

**Background:** Phytoestrogen is a plant-derived compound, which has estrogenic effect and it is found in liquorice root extract and sesame oil. **Aim of the work:** To investigate some biochemical effects of liquorice root extract and sesame oil on male albino rats.

**Materials and methods:** 18 animals were divided randomly into three groups. **Group A:** Control group, **group B:** rats treated with oral dose of liquorice 10 mg/kg body weight/day for one month, and **group C:** rats treated with oral dose of sesame oil 1 ml/kg body weight/day. At the end of the experiment, blood samples were collected for biochemical analysis.

**Results:** Liquorice and sesame oil induced highly significant decrease in TC, TG, LDL, VLDL, LDL/HDL ( $p < 0.01$ ) and significant increase in HDL ( $p < 0.05$ ). They also showed highly significantly decrease in FSH, testosterone and sperm count compared to control group ( $p < 0.01$ ).

**Conclusion:** The study supports that the high level intake of liquorice root extract or sesame oil caused hormonal disturbance and decreases sperm count.

**Keywords:** Liquorice root, Sesame oil, Lipid profile, Albino rats, Physiological parameters.

### INTRODUCTION

*Glycyrrhizaglabra*, a family of Leguminosae, is a plant that grows in Egypt and other countries of the world. Its roots possess some nutritional value and medicinal properties. It is widely used as a cold drink and in the preparation of some pharmaceuticals such as hematinic pills<sup>(1)</sup>.

Phytochemical analysis of *Glycyrrhizaglabra* (liquorice) root extract showed that it contains saponin triterpenes (glycyrrhizin, glycyrrhetic acid and liquiritic acid), flavonoids (liquiritin, isoflavonoids and formononetin) and other constituents such as coumarins, sugars, amino acids, tannins, starch, choline, phytosterols and bitter principles, which are most likely responsible for its therapeutic properties<sup>(2)</sup>.

There are many phenols present in liquorice extracts such as liquiritigenin, liquiritin, isoliquiritigenin, isoliquiritin, glabridin and formononetin, which are responsible for the plant estrogen activity of this plant, extracts<sup>(3)</sup>.

Medicinally its root is used widely either as tincture or fluid extract or ingredient in over the counter pharmaceutical products for cough, expectorant, asthma, peptic ulcers and hepatic-protective products. This is because of its anti-inflammatory, anti-ulcerative demulcent, antimicrobial and hepatic-protective activities<sup>(4)</sup>. In Japan, intravenously, liquorice components are used for treating hepatitis B and C (National Center for Complementary and Integrative Health, 2016). Other

indications still under research like different cancer conditions (Colon, Breast, Hepatic and Prostate),

anti-diabetic, antiangiogenic, neuro-protective, rheumatoid arthritis and cardio-protective<sup>(4)</sup>.

### Sesame oil

Sesame belongs to the family Pedaliaceae. It is one of the richest food source of lignans, a major type of phytoestrogens known to man since the dawn of civilization<sup>(5)</sup> and is increasingly incorporated into human diets because of its health benefits<sup>(6)</sup>. Sesame lignans such as sesamin, episesamin, sesamol, and sesamol isolated from *Sesamum indicum* seeds are implicated as having certain properties such as anti-tumorigenic<sup>(7)</sup> and antioxidant<sup>(8)</sup>. The proximate analysis of sesame seed indicates that it contains about 50-60 % oil, 8 % protein, 5.8 % water, 3.2 % crude fiber, 18 % carbohydrate and 5.7 % ash. It is very rich in minerals such as calcium, phosphorus and vitamin E<sup>(9)</sup>. The seed oil combats some health conditions like cold and chronic cough and in turn prevents bronchial lung disease. In addition, it helps in improving the blood glucose, glycosylated hemoglobin and lipid-peroxidation<sup>(10)</sup>. Besides its broad use in cooking, the oil is also used in the manufacture of margarine and some pharmaceuticals<sup>(11)</sup>. Sesamin, the most abundant lignan in sesame oil<sup>(12)</sup> is known to contain unique amounts of phytoestrogen, and has some effects on sex physiology<sup>(12)</sup>.

## MATERIALS AND METHODS

Roots of liquorice plant were purchased from market of Agricultural Seeds and Medicinal Plants, Cairo, Egypt. Clean roots (10gm) were soaked in cup of water over night and then filtered. This extract was freshly prepared every day.

Sesame oil extracts were purchased from Cap Pharm for Extracting Natural Oils and Herbs, Cairo, Egypt.

### Experimental animal

The experiment was carried out on 18 male albino rats *Rattusrattus* strain weighing (130-140gm) obtained from animal farm of El-Nile Company for Pharmaceutical Products, Cairo, Egypt. Animals were housed in metallic cages and maintained under standard condition of temperature, humidity and natural light/dark cycle along the experimental period. Food and water were available throughout the experimental *ad libitum*. Rats were left to acclimatize for one week before starting the experiment.

### Experimental design

Rats were divided into three equal groups (6 rats in each group) as the following:

**Group I** (control group): maintained on standard pellet diet and tap water *ad libitum* for 30 days.

**Group 2:** rats received orally sesame oil (1 ml/kg body weight/day) for 30 days.

**Group 3:** rats received orally liquorice (*Glycyrrhizaglabra*) root aqua extract (10 mg/kg body weight/day) for 30 days.

### Body weight measurement

Body weight was recorded on zero time and at the end of the experiment.

### Blood sample collection

At the end of the experimental period, the blood samples were collected from the retro-orbital sinus after overnight fasting and the rats were anesthetized by ether. Serum was separated by centrifugation at 2500 g for 15 minutes at room temperature to estimate biochemical parameters.

### Biochemical analysis

#### Assessment of biochemical parameters:

In the present study, total protein (TP) and albumin concentration were estimated, then serum globulin concentrations were calculated according to the formula: Globulin (g/dl) = total protein (g/dl) – albumin (g/dl). Aspartate aminotransferase (AST), alanine aminotransferase (ALT), creatinine, BUN

and blood glucose concentrations as well as lipid profile including total cholesterol, triglycerides and high-density lipoprotein cholesterol (HDL-C) were also determined. All parameters were estimated using BioMerieux SA kits, France.

Albumin/globulin ratio was determined. In addition, ratios of TC/HDL (risk factor 1) and LDL/HDL (risk factor 2) were also calculated after calculation of serum LDL-C (low-density lipoprotein cholesterol) and VLDL (very low density lipoprotein cholesterol) using the Friedewald's<sup>(13)</sup> and Norbert<sup>(14)</sup> formulas, respectively as following:

Friedewald's equation:

$$\text{LDL (mg/dl)} = \text{TC} - \{\text{HDL} + [\text{TG}/5]\}.$$

Norbert equation:  $\text{VLDL} = \text{TG}/5$ .

TC/HDL (risk factor 1)

LDL/HDL (risk factor 2)

### Hormonal assay

Estimation of serum luteinizing hormone (LH), follicle-stimulating hormone (FSH) and testosterone (T) levels by following manufacture instructions of kit. All kits used for hormone assay were Monobind Inc., Lake Forest CA 92630, USA).

### Sperm collection and evaluation

The left caudal epididymis was separated and the total recovered sperm during 4 h of incubation in normal saline (volume = 1 ml, 35~37°C) was calculated. The sperm concentration was determined by the conventional method using a hemocytometer chamber for the red blood cell count<sup>(15)</sup>.

### Ethical approved

**The study was approved by the Ethics Board of Al-Azhar University.**

### Statistical analysis

The results were expressed as mean ± SEM (standard error of the mean). Data were analyzed by T-test and were performed using the Statistical Package (SPSS) program, version 19. The Bonferroni test was used as a method to compare significance between groups.

## RESULTS

### Body weight and glucose level

Results of the present study showed a significant increase ( $p < 0.05$ ) in body weight in the treated groups and non-significant change in glucose level in the treated groups when compared to control rats (Table 1).

**Table(1):**Percentagechanges in body weight and glucose levelin control, sesame oil and liquorice groups.

Group Parameters	Control	Liquorice	Sesame oil
Body weight	137.6 ± 0.4	190.6±0.5* 38.5%	195.1±0.4* 41.8%
FBS (mg/dl)	94.0±2.8	88.3±2.0 -6.6%	90±1.9 -4.2%

Values represent mean ±SEM. (p\* $<$ 0.05, as compared to control group).

#### Kidney functions

The data in table (2) showed non-significantchange in urea and serum creatinine in liquoriceand sesame groups as compared to control (Table 2).

**Table(2):** Changes in the BUN and creatinine levels in the control,liquoriceand sesame groups.

Groups Parameters	Control	Liquorice Extract	Sesame oil
BUN (mg/dl)	20.3±0.2	16.6±0.4 -18.2%	20.1±0.2 -0.9%
Creatinine (mg/dl)	0.7±0.2	0.6±0.01 -28.5%	0.6±0.001 14.3%

Values represent mean ±SEM.

#### Liver functions

Results of the present study showednon-significantchange in ALAT and ASAT in the liquoriceand sesame groups when compared to control rats (Table 3).

**Table (3):** Changes in ALAT and ASAT in the control, liquoriceand sesame groups.

Groups Parameters	Control	Liquorice extract	Sesame oil
ALAT(U/L)	22.5±0.6	20.0±0.4 -11.1%	18.9±0.3 16%
ASAT(U/L)	33.5±0.5	29.1±0.3 -13.1%	32.1±0.2 4.2%

Values represent mean ±SE (stander error).

#### Lipid profile

Results of the present study showed that liquoriceand sesame produced highly significant decrease (p $<$ 0.01) in TC,TG, LDL, VLDL and LDL/HDLwhen compared to control group.While, there was highly significant increase (p $<$ 0.05) in HDL, as compared to control animals (Table 4).

**Table (4):** Changes in the lipid profile in the control, liquorice andsesame groups.

Groups Parameters	Control	Liquorice extract	Sesame oil
T C(mg/dl)	141.3±0.42	86.3±1.5** -38.9%	85±0.01** -39.8%
T G(mg/dl)	71.5±1.56	56.8±4.4* -20.5%	43.5±2.1** -39.1%
HDL(mg/dl)	50.3±0.33	61.3±0.33* 21.8%	60±0.01* 19.3%
LDL(mg/dl)	71.2±1.27	14.1±1.1** -80.1%	16.3±.58** -77.1%
VLDL(mg/dl)	14.3±0.31	11.0±0.9* -23.0%	8.7±0.58** -39.1%
LDL/HDL(mg/dl)	1.22±0.007	0.22±.01** -81.9%	0.27±0.01** -77.8%
TC/HDL(mg/dl)	2.4±0.008	1.4±.22** -41.6%	1.4±0.01** 41.6%

Values represent mean ±SEM. (p\* $<$ 0.05, p\*\* $<$ 0.001 as compared to control group).

### Protein profile

In the present study, administration of liquorice and sesame to normal rats showed non-significant change in total protein, albumin, globulin and albumin/globulin ratio in the treated groups when compared to control rats (Table 5).

**Table (5):** Changes in the total protein, albumin, globulin and albumin/globulin levels in control and treated groups.

Groups Parameters	Control	Liquorice extract	Sesame oil
Total protein (g/dl) % of change	6.4±0.12	5.5±0.20 -14.0%	6.0±0.1 -6.25%
Albumin (g/dl) % of change	3.7±0.11	4.0±.15 8.1%	3.5±0.19 -5.4%
Globulin (g/dl) % of change	2.7±0.10	2.3±0.22 -14.8%	2.5±0.3 -7.4%
Albumin/Globulin(g/dl) % of change	1.3±0.1	1.7±0.06 30.7%	1.4±0.06 7.6%

Values represent mean ±SEM.

### Hormones

The present study showed that administration of liquorice extract to normal rats showed a highly significant decreased FSH and testosterone ( $p < 0.01$ ), while in sesame group there were a significantly decreased FSH and testosterone ( $p < 0.05$ ) when compared to control rats (Table 6).

**Table(6):** Changes in FSH, LH and Testosterone levels in control, sesame and liquorice groups.

Groups Parameters	Control	Liquorice extract	Sesame oil
FSH(ng/ml) % of change	2.9±0.1	1.9±0.1** -34.4%	2.0±0.1* -31.0%
LH(ng/ml) % of change	1.9±0.1	1.7±0.1 -10.5%	1.8±0.1 -5.2%
Testosterone(µ /dl) % of change	3.9±0.2	2.8±0.2** -39.3%	2.9±0.2* -25.06%

Values represent mean ±SE (stander error). ( $p^* < 0.05$ ,  $p^{**} < 0.001$  as compared to control group)

### Sperm count

The data in table (7) showed decrease in sperm count in liquorice and sesame groups when compared to control rats.

**Table (7):** Sperm count in the control, sesame and liquorice group.

Groups Parameters	Control	Liquorice extract	Sesame oil
Sperm count*10 <sup>6</sup> /ml % of change	100	38±0.01 -62%	63±0.01 -37%

## DISCUSSION

### Body weight

The findings in this study showed an increase in body weight in both experimental groups treated with *Glycyrrhizaglabra* (liquorice) and sesame oil. In *Glycyrrhizaglabra* the increase in body weight by *Glycyrrhizaglabra* root extract is in accordance with that reported by Miller<sup>(16)</sup> who demonstrated that *Glycyrrhizaglabra* inhibits 11 β-hydroxysteroid dehydrogenase and induces excess release of mineralocorticoids, which causes retention of sodium

and water that leads to edema and increase in body weight. However, the increased body weight is explained by the improvement in feeding efficiency<sup>(17)</sup>.

While in Sesame oil group the increase in body weight was due to unsaturated fatty acids such as oleic acid and linoleic acid, which increase the effectiveness of non-steroidal hormones such as thyroid and growth hormones which are important

factors in increasing body weight by stimulating the basal metabolic rate<sup>(18)</sup>.

### Lipid profile

The decrease in serum total cholesterol and triglycerides reported in this study following oral administration of *Glycyrrhiza glabra* (liquorice) root for 4 weeks is similar to that reported by **Khushbaktova *et al.***<sup>(19)</sup> and **Fuhrman *et al.***<sup>(20)</sup>. The studies of previously mentioned authors attributed the hypocholesterolemic effect of liquorice to the presence of certain isoflavones, which act as antioxidants via inhibition of LDL oxidation which inhibits the local mechanism of atherogenesis. Moreover, **Nikitina *et al.***<sup>(21)</sup> reported that the glycosides of *Glycyrrhiza glabra* prevent accumulation of cholesterol in cells as well as human blood serum. A significant decline in plasma LDL-cholesterol in treated groups could be correlated with saponin content of liquorice root, whereas saponin enhances the hepatic LDL-receptor levels, increase hepatic uptake of LDL-cholesterol and aids its catabolism to bile acid<sup>(22)</sup>. Saponins are known to lower triglyceride by inhibiting pancreatic lipase activity. While, the decline in VLDL cholesterol levels in the treated groups could be directly correlated to decline in triglyceride levels of these groups, as it is well established that VLDL particles are the main transporters of triglyceride in plasma<sup>(23)</sup>. Phytosterols are reported to displace intestinal cholesterol and reduce cholesterol absorption from intestine<sup>(24)</sup>. Saponins are capable of precipitating cholesterol from micelles and interfere with enterohepatic circulation of bile acids, making it unavailable for intestinal absorption<sup>(25)</sup>.

In sesame oil group, there was a significant decrease in lipid profile. Sesame lignans (sesamin and/or episesamin) reduced serum and liver cholesterol concentrations by inhibiting absorption and synthesis of cholesterol. **Ogawa *et al.***<sup>(26)</sup> reported that daily oral intake of sesamin in hypercholesterolemic patients for 4 weeks significantly decreased total cholesterol and LDL-C concentrations. In rat, the mechanism for the hypocholesterolemic effect of sesamin is believed to be related to inhibition of intestinal absorption of cholesterol, increased excretion of cholesterol into bile and decreased activity of 3-hydroxy-3-methylglutaryl coenzyme A reductase.

### Hormones

Change in hormonal levels in this study showed a significant reduction in follicle stimulating hormone and testosterone. Testosterone is a steroid hormone from androgen group, and is one of the major sex hormones produced by the body in both men and women. It plays key roles in health and

well-being. FSH and testosterone are involved as synergistic in the process of spermatogenesis<sup>(27)</sup>. **Armanini *et al.***<sup>(28)</sup> found that liquorice reduces serum testosterone in healthy men. Also liquorice inhibits 6  $\beta$ -hydroxy steroid dehydrogenase and stimulating aromatase, resulting in reduced serum testosterone. Liquorice reduced the serum testosterone level and affected androgen metabolism by inhibiting the enzyme  $\beta$ -6 HSD and 17- $\beta$ HSD hydroxysteroid dehydrogenase or by stimulating aromatase. Therefore, it was proposed that liquorice causes the deficiency in serum testosterone<sup>(29)</sup>, glabridin, the major isoflavone in liquorice and many other phenols such as liquiritigenin, liquiritin, isoliquiritigenin, isoliquiritin, glabridin and formononetin that are responsible for phytoestrogen activity and estrogenic properties<sup>(30)</sup>, which cause hormonal disturbance and decrease the level of FSH and Testosterone, which lead to decrease in sperm count.

While in sesame oil group, the decrease in serum FSH and testosterone was approved by the study of **Brown and Chakraborty**<sup>(31)</sup> who showed that estrogen agonist like clomiphene has been found to decrease the synthesis and/or release of gonadotrophins with the implication of both low serum LH and testosterone concentration found in male rats. FSH concentration decreased in a high dose of sesame group compared to control group. **Shittuet *et al.***<sup>(32)</sup> found that sesame phytoestrogen lignans can stimulate testosterone aromatization to estradiol, or convert it to Dihydrotestosterone.

### CONCLUSION

Therefore, it can be concluded that the decrease in the concentrations of testosterone was due to its conversion into estradiol which is done by aromatase and reductase enzymes and sesame lignans. This study supposes that the decrease in FSH and testosterone level leads to decreasing in the sperm count.

### REFERENCE

- Fenwick G, Lutomski J, Nieman C (1990):** *Glycyrrhiza glabra L.* (Liquorice): Composition, uses and analysis. *Food Chem.*, 38(2): 119-143.
- Arystanova T, Irismetov M, Sophekova A (2001):** Chromatographic determination of glycyrrhizic acid in *Glycyrrhiza glabra* preparation. *Chem Nat Com.*, 37: 89-91
- Tang Z L, Tong Y, Chen X, Chen X, Wang Y, Lu J, Angeles L (2015):** A systematic review of the anticancer properties of compounds isolated from licorice (Gancao). *Planta Med.*, 18: 1670–1687.
- Peng F, Du Q, Peng C, Wang N, Tang H, Xie X, Shen J, Chen J (2015):** A review: The pharmacology of Isoliquiritigenin. *Phytotherapy Research*, 29:969–977.

5. **Zavareh M, Hoogenboom G, Rahimian M H, Arab D A (2008):** A decimal code to describe the growth stages of sesame (*Sesamum orientale* L.). *Int J Plant Prod.*, 2:193-206.
6. **Shittu L A J, Shittu R K, Adesite S O, Ajala M O, Bankole, M A, Benebo AS (2008):** *Sesame radiatum* Phytoestrogens Stimulate Spermatogenic Activity and Improve Sperm Quality in Adult Male Sprague Dawley Rat Testis. *Int J Morphol.*, 26(3):643-652.
7. **Hirose N, Inoue T, Nishihara K, Sugano M, Akimoto K, Shimizu S, Yamada H (1991):** Inhibition of cholesterol absorption and synthesis in rats by sesame. *J. Lipid Res.*, 32:629-638.
8. **Jeng KCG, Hou RCW (2005):** Sesamin and Sesamol: Nature's Therapeutic Lignans. *Curr Enzym Inhib.*, 1:11-20.
9. **Obiajunwa EI, Adebisi FM, Omode PE (2005):** Determination of Essential Minerals and Trace Elements in Nigerian Sesame Seeds, Using TXRF Technique. *Pakistan J Nutr.*, 4: 393-395.
10. **Kanu PJ (2011):** Biochemical analysis of black and white sesame seeds from China. *Am J Biochem Mol Biol.*, 1:145-157.
11. **Sangha M, Gupta P, Tharpo V (2004):** Storage studies on plant oils and their methyl esters. *E Journal*, 7:1-15.
12. **Fukuda Y, Nagata M, Namiki MJ (1986):** Contribution of Lignan Analogues to Antioxidant Activity of Refined Unroasted Sesame Seed Oil. *J Am Oil Chem. Soc.*, 63:1027-1031.
13. **Jacklin A, Ratledge C, Welham K, Bilko D, Newton CJ (2003):** The sesame oil constituent, sesamol, induces growth arrest and apoptosis of cancer and cardiovascular cells. *Ann. New York Acad Sci.*, 1010:374-380.
14. **Friedwald's WT, Levy RI, Fredrickson DS et al. (1999):** Estimation of the concentration of low-density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge. *Clin Chem.*, 18:499-502.
15. **Norbert WT (1995):** clinical guide to laboratory tested. 3<sup>rd</sup> ed. Saunders W.B., Company, Philadelphia.
16. **WHO (2010):** WHO laboratory manual for the examination and processing of human semen. Fifth ed. Geneva, Switzerland: WHO press: 22-44. <https://apps.who.int/iris/handle/10665/44261>
17. **Miller I (1998):** Herbal medicine: Selected clinical considerations focusing on known or potential drug-herb interaction. *Arch. Inter Med.*, 158: 2200-2211.
18. **Alaauldeen S M AL-Sallami (2017):** Effect of Sesame Oil on Male Rats Treated with Acrylamide in some Physiological and Hormonal Blood Criteria. *IJCPR.*, 8: 134-140.
19. **Khushbaktova Z, Syrov V, Batirov E (1991):** Effect of flavonoids on the course of hyperlipidemia and atherosclerosis under experimental conditions. *Khimiko Farmatsevticheskii Zhurnal*, 25:54- 57.
20. **Fuhrman B, Vaya J, Belinky P, Aviram M (1999):** The isoflavone glabridin inhibits LDL oxidation. <https://link.springer.com/content/pdf/10.1007%2F3-540-27661-0.pdf>
21. **Nikitina N, Khalilov E, Tertov V, Orekhov A (1995):** *In vitro* decrease in the atherogenicity by polyunsaturated phosphotidylcholine. *Byu. Eksper. Biol Med.*, 119: 497-501.
22. **Harwood J H J, Chandler C E, Pellarin L D, Bangerter F W, Wilkin RW, Long CA (1993):** Pharmacologic consequences of cholesterol absorption inhibition: alteration in cholesterol metabolism and reduction in plasma cholesterol concentration induced by the synthetic saponin betatigogenin cellobioside (CP-88818; tiqueside). *J Lipid Res.*, 34: 377-95.
23. **Howell T J, MacDougall D E, Jones P J H (1998):** Phytosterols partially explain differences in cholesterol metabolism caused by corn or olive oil feeding. *J Lipid Res.*, 39: 892-900.
24. **Ikeda I, Sugano M (1998):** Inhibition of cholesterol absorption by plant sterols for mass intervention. *Curr Opin Lipidol.*, 9: 527-531.
25. **Oakenfull D G, Sidhu G S (1990):** Could saponins be a useful treatment for hypercholesterolaemia? *Eur J Clin Nutr.*, 44: 79-88
26. **Ogawa H, Sasagawa S, Murakami T, Yoshizumi H (1995):** Sesame lignans modulate cholesterol metabolism in the stroke-prone spontaneously hypertensive rat. *Clin Exp Pharmacol Physiol.*, 22: 310-312.
27. **Berne RM, Levy MN (1993):** Physiology. Third ed. St. Louis, Mosby, pp: 950-959.
28. **Armanini D, Bonanni G, Mattarello M J, Fiore C, Sartorato P, Palermo M (2003):** Licorice Consumption and Serum Testosterone in Healthy Man. *Endocrinol Diabetes Exp Clin.*, 111: 341-343.
29. **Tamir S, Eizenberg M, Somjen D, Stern N, Shelach R, Kaye A, Vaya J (2000):** Estrogenic and anti-proliferative properties of glabridin from licorice in human breast cancer cells. *Cancer Res.*, 60: 5704-5709.
30. **Tang Z, Li T, Tong Y, Chen X, Chen X, Wang Y, Lu J, Angeles L (2015):** A systematic review of the anti-cancer properties of compounds isolated from licorice (Gancao). *Planta Med.*, 81:1670-1687.
31. **Brown JL, Chakraborty PK (1998):** Characterization of the effects of clomiphene citrate on reproductive physiology in male rats of various ages. *Acta Endocrinol.*, 118:437-443.
32. **Shittu LAJ, Shittu RK, Adesite SO, Ajala MO, Bankole MA, Benebo AS (2008):** Sesame radiatum Phytoestrogens Stimulate Spermatogenic Activity and Improve Sperm Quality in Adult Male Sprague Dawley Rat Testis. *Int J Morphol.*, 26:643-652.